Recommended within 72 hours of assault

Please consider all risks, benefits, and contraindications before the initiation of therapy.

Risk Category	Clinical/Historical Findings	Recommendation for Non-Occupational Post Exposure Prophylaxis	Medication Recommendations
High	Penetration assault by one or more assailants known to be HIV positive or at high risk of HIV infection (Injection drug users, men who have sex with men)* and / or Anal penetration with or without injuries ** *HIV virus prevalence among convicted sex offenders may be twice the general male population which emphasizes the higher risk of HIV exposure following sexual assault **Injuries include trauma / tearing of mucosal tissue and bleeding or presence of blood	Strongly Recommended	Truvada, (Emtracitabine, 200 mg/ Tenofovir 300 mg), 1 tablet po daily x 28 days and Isentress (Raltegravir 400 mg), 1 tablet po twice daily x 28 days
Moderately High	Penetration assault by one or more assailants of unknown HIV status with vaginal injuries and known or unsure ejaculation and no or uncertain condom use Penetration assault in the presence of sexually transmitted infection, genital lesion, IUD, menstruation Penetration assault by one or more uncircumcised assailants of unknown HIV status	Recommended	As per high risk
Moderately Low	Penetration assault by one or more assailants of unknown HIV status with no vaginal injuries with ejaculation or vaginal injuries without ejaculation	Optional	As per high risk
Low	No anal or vaginal penetration No ejaculation from the assailant Oral penetration only Condom use Assailant known to be HIV negative Bite injury unless the biter's mouth was bloody and the exposed patient's skin is visibly broken	Not Recommended	None
N/A	Assault occurred greater than 72 hours ago		Consult Infectious Disease or NPEP Hotline 1-888-448-4911 as needed